






















Infants', Children's and Jr. TYLENOL®

Use this chart to determine the proper dose of TYLENOL® for your child.

mL = milliliter
tsp = teaspoonful

If possible, use weight to dose; otherwise use age

DOSE: Every 4 hours as needed. DO NOT GIVE MORE THAN 5 DOSES IN 24 HOURS.

Weight (lbs)	Age	 Infants' TYLENOL® Oral Suspension Active Ingredient: Acetaminophen 160mg (in each 5mL or 1 tsp) Use only as directed.	 Children's TYLENOL® Oral Suspension Active Ingredient: Acetaminophen 160mg (in each 5mL or 1 tsp) Use only as directed.	 Children's TYLENOL® Meltaway Chewable Tablets Active Ingredient: Acetaminophen 80mg (in each tablet) Use only as directed.	 Jr. TYLENOL® Meltaway Chewable Tablets Active Ingredient: Acetaminophen 160mg (in each tablet) Use only as directed.
6-11 lbs	0-3 mos	1.25 mL 	—	—	—
12-17 lbs	4-11 mos	2.5 mL 	—	—	—
18-23 lbs	12-23 mos	3.75 mL 	—	—	—
24-35 lbs	2-3 yrs	5 mL 	5 mL (1 tsp) 	2 tablets 	—
36-47 lbs	4-5 yrs	—	7.5 mL (1½ tsp) 	3 tablets 	—
48-59 lbs	6-8 yrs	—	10 mL (2 tsp) 	4 tablets 	2 tablets 
60-71 lbs	9-10 yrs	—	12.5 mL (2½ tsp) 	5 tablets 	2½ tablets 
72-95 lbs	11 yrs	—	15 mL (3 tsp) 	6 tablets 	3 tablets 

- Take every 4 hours as needed
- Do NOT exceed more than 5 doses in 24 hours
- Do NOT use with any other product containing acetaminophen
- Keep all medicines out of the reach of children
- Do NOT administer adult medicines to children

Infants' and Children's Motrin®

Use this chart to determine the proper dose of Motrin® for your child.

mL = milliliter
tsp = teaspoonful

If possible, use
weight to dose;
otherwise use
age

DOSE: Every 6–8 hours as needed. DO NOT GIVE MORE THAN 4 DOSES IN 24 HOURS.



Infants' MOTRIN® Concentrated Drops

Active Ingredient:
Ibuprofen 50 mg (NSAID)*
(in each 1.25 mL)

*Nonsteroidal
anti-inflammatory drug
Use only as directed.

Available in:
 Dye-Free
Berry



Children's MOTRIN® Oral Suspension

Active Ingredient:
Ibuprofen 100 mg (NSAID)*
(in each 5 mL or 1 tsp)

*Nonsteroidal
anti-inflammatory drug
Use only as directed.

Available in:
 Original
Berry
 Dye-Free
Berry

WEIGHT	AGE		
6-11 lbs	0-5 mos	Do not use	—
12-17 lbs	6-11 mos	1.25 mL	
18-23 lbs	12-23 mos	1.875 mL	
24-35 lbs	2-3 yrs	—	5 mL (1 tsp)
36-47 lbs	4-5 yrs	—	7.5 mL (1½ tsp)
48-59 lbs	6-8 yrs	—	10 mL (2 tsp)
60-71 lbs	9-10 yrs	—	12.5 mL (2½ tsp)
72-95 lbs	11 yrs	—	15 mL (3 tsp)

- Take every 6–8 hours as needed
- Do NOT exceed more than 4 doses in 24 hours
- Do NOT administer longer than 10 days, unless directed by a doctor
- Ask a doctor or pharmacist before use if the child is taking any other drug containing an NSAID (prescription or non prescription)